

<b>REPORT DOCUMENTATION PAGE</b>			Form Approved OMB NO. 0704-0188		
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA, 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 14-02-2016		2. REPORT TYPE Final Report		3. DATES COVERED (From - To) 1-Jun-2011 - 31-Aug-2014	
4. TITLE AND SUBTITLE Final Report: Fractional Differential and Integral Inequalities with Applications			5a. CONTRACT NUMBER W911NF-11-1-0047		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER 611102		
6. AUTHORS Donna Stutson, A.S. Vatsala			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAMES AND ADDRESSES Xavier University of Louisiana 1 Drexel Drive  New Orleans, LA 70125 -1098			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS (ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			10. SPONSOR/MONITOR'S ACRONYM(S) ARO		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) 57873-MA-H.11		
12. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.					
14. ABSTRACT The monotone method extended to such systems is called the generalized monotone method. Here the generalized monotone method has been extended to the Caputo fractional differential equation of order $q$ (where $0 < q < 1$ ) with an initial condition as well as the existence of coupled minimal and maximal solutions for such an equation and a numerical example is provided as an application of the theoretical results.  <del>The linear sub-hyperbolic fractional partial differential equation in one dimensional space, that is the <math>n</math>th time order</del>					
15. SUBJECT TERMS Caputo Derivative, Caputo Partial Differential Equations, Eigenfunctions, Upper and Lower Solutions, Generalized Monotone Method					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	15. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT UU	b. ABSTRACT UU	c. THIS PAGE UU			Donna Stutson
					19b. TELEPHONE NUMBER 504-520-5308



## Report Title

Final Report: Fractional Differential and Integral Inequalities with Applications

### ABSTRACT

The monotone method extended to such systems is called the generalized monotone method. Here the generalized monotone method has been extended to the Caputo fractional differential equation of order  $q$  (where  $0 < q < 1$ ) with an initial condition as well as the existence of coupled minimal and maximal solutions for such an equation and a numerical example is provided as an application of the theoretical results.

The linear sub hyperbolic fractional partial differential equation in one dimensional space, that is the  $q$ th time order derivative is such that  $1 < q < 2$ , and the linear super hyperbolic fractional partial differential equation in one dimensional space, that is the  $q$ th time order derivative is such that  $2 < q < 3$  are considered. In the special case when  $q = 1$ , it is the parabolic equation and when  $q = 2$ , it is the hyperbolic equation. The eigenfunction expansion method is used to obtain representation forms for the solution of the linear sub and the linear super hyperbolic fractional partial differential equations in terms of the nonhomogeneous terms, initial and boundary conditions. See [9,10, 11, 12, 14] and the references therein for qualitative study, applications and numerical study of fractional differential equations. The representation forms obtained are useful in computing numerical solutions to the sub and super hyperbolic partial differential equations.

---

**Enter List of papers submitted or published that acknowledge ARO support from the start of the project to the date of this printing. List the papers, including journal references, in the following categories:**

**(a) Papers published in peer-reviewed journals (N/A for none)**

Received

Paper

**TOTAL:**

**Number of Papers published in peer-reviewed journals:**

---

**(b) Papers published in non-peer-reviewed journals (N/A for none)**

Received

Paper

**TOTAL:**

**Number of Papers published in non peer-reviewed journals:**

---

**(c) Presentations**

"Generalized Monotone Method, Numerical Approach for Ordinary Fractional Differential Equations." by Aghalaya S. Vatsala, ULL, Sowmya Muniswamy, ULL and Donna Stutson, Xavier University of Louisiana (1083-34-93) at The Fall Southeastern section Meeting at Tulane University, New Orleans, LA, October 13-14, 2012, meeting # 1083.

"A Representation Formula for the One Dimensional Caputo Fractional Reaction Diffusion Equation and a Numerical Example Using the Generalized Monotone Method." by Donna Stutson, Xavier University of Louisiana and Aghalaya S. Vatsala, ULL,(1083-35-163)at The Fall Southeastern section Meeting at Tulane University, New Orleans, LA, October 13-14, 2012, meeting # 1083.

"Numerical Methods for Fractional Differential Equations Via Generalized Monotone Method." by Aghalaya S. Vatsala, ULL, Sowmya Muniswamy, ULL and Donna Stutson, Xavier University of Louisiana (1086-34-1007) at the Joint Mathematics Meeting , January 9-12, 2013, San Diego Convention Center.

"A Numerical Example for the One Dimensional Caputo Fractional Wave Equation Using the Representative Solution and the Generalized Monotone Method." by Donna Stutson, Xavier University of Louisiana and Aghalaya S. Vatsala, ULL (1086-35-2418) at the Joint Mathematics Meeting , January 9-12, 2013, San Diego Convention Center.

"Riemann Liouville and Caputo Fractional Differential and Integral Inequalities with Applications." by Aghalaya S. Vatsala, ULL, Sowmya Muniswamy, ULL and Donna Stutson, Xavier University of Louisiana at New Trends in Differential Equations, Department of Mathematics, University of Tennessee at Chattanooga, Chattanooga, TN 37403-2598, March 15-17, 2013.

Superlinear Convergence via Iterative Methods for Scalar Caputo fractional differential equations with applications.  
Sowmya Muniswamy\*, University of Louisiana at Lafayette  
Aghalaya S. Vatsala, University of Louisiana at Lafayette  
(1096-34-1660)

Super Hyperbolic Linear Partial Fractional Differential Equations in One Dimensional Space. D S Stutson\*, Xavier University of Louisiana  
A S Vatsala, University of Louisiana at Lafayette  
(1096-35-2439)

Sub and Super Hyperbolic Linear Partial Fractional Differential Equations with Numerical Results.  
Aghalaya S. Vatsala\*, University of Louisiana at Lafayette  
Donna Sue Stutson, Xavier University of Louisiana  
(1096-35-1328)

**Number of Presentations: 8.00**

---

**Non Peer-Reviewed Conference Proceeding publications (other than abstracts):**

Received

Paper

**TOTAL:**

**Number of Non Peer-Reviewed Conference Proceeding publications (other than abstracts):**

---

**Peer-Reviewed Conference Proceeding publications (other than abstracts):**

Received

Paper

08/03/2012 1.00 Donna Stutson, A.s. Vatsala. A representative solution is obtained for the onedimensional Caputo Fractional Reaction DiffusionEquation,  
Sixth International Conference on Dynamical Systems and Applications. 27-MAY-11, . : ,

**TOTAL: 1**

**(d) Manuscripts**

<u>Received</u>	<u>Paper</u>
01/08/2016 9.00	TH. T PHAM, J. D. RAMIREZ, A. S. VATSALA. GENERALIZED MONOTONE METHOD FOR CAPUTO FRACTIONAL DIFFERENTIAL EQUATION WITH APPLICATIONS TO POPULATION MODELS, Neural, Parallel and Scientific Computations (03 2013)
01/08/2016 8.00	Donna S Stutson, Aghalaya S. Vatsala. Super Hyperbolic Linear Partial Fractional Differential Equation in One Dimensional Space, Communications in Applied Analysis (01 2014)
01/11/2014 7.00	DONNA STUTSON,AGHALAYA S VATSALA. RIEMANN LIOVILLE AND CAPUTO FRACTIONALDIFFERENTIAL AND INTEGRAL INEQUALITIES, Dynamic Systems and Applications (07 2013)
02/14/2016 10.00	Aghalaya S. Vatsala, Bhuvaneswari Sambandham. Laplace Transform Method for Sequential CaputoFractional Differential Equations / MESA, Nonlinear Studies (08 2015)
08/10/2012 2.00	Donna Stutson, A. S. Vatsala. GENERALIZED MONOTONE METHOD FORFRACTIONAL REACTION DIFFUSION EQUATIONS, Communications in Applied Analysis (09 2011)
08/10/2012 3.00	A.S. VATSALA, TH.T PHAM, J.D. RAM!!IREZ. GENERALIZED MONOTONE METHOD FOR CAPUTOFRACTIONAL DIFFERENTIAL EQUATION WITHAPPLICATIONS TO POPULATION MODELS, Neural, Parallel & Scientific Computations (04 2012)
08/24/2013 4.00	SOWMYA MUNISWAMY, AGHALAYA S VATSALA. NUMERICAL APPROACH VIA GENERALIZED MONOTONEMETHOD FOR SCALAR CAPUTOFRACTIONAL DIFFERENTIAL EQUATIONS, Neural, Parallel & Scientific Computations (01 2013)
08/24/2013 5.00	SOWMYA MUNISWAMY, AGHALAYA S. VATSALA. SUPERLINEAR CONVERGENCE FOR CAPUTO FRACTIONALDIFFERENTIAL EQUATIONS WITH APPLICATIONS, Dynamic Systems and Applications (04 2013)
08/24/2013 6.00	Donna S. Stutson, Aghalaya S Vatsala. Sub Hyperbolic Linear Partial FractionalDifferential Equation in One Dimensional Spacewith Numerical Results, Neural, Parallel & Scientific Computations (07 2013)
<b>TOTAL:</b>	<b>9</b>

Number of Manuscripts:

Books

Received      Book

TOTAL:

Received      Book Chapter

TOTAL:

Patents Submitted

Patents Awarded

Awards

Graduate Students

<u>NAME</u>	<u>PERCENT_SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Names of Post Doctorates

<u>NAME</u>	<u>PERCENT_SUPPORTED</u>
FTE Equivalent:	
Total Number:	

---

### Names of Faculty Supported

NAME

PERCENT SUPPORTED

**FTE Equivalent:**

**Total Number:**

---

### Names of Under Graduate students supported

NAME

PERCENT SUPPORTED

**FTE Equivalent:**

**Total Number:**

---

### Student Metrics

This section only applies to graduating undergraduates supported by this agreement in this reporting period

The number of undergraduates funded by this agreement who graduated during this period: .....

The number of undergraduates funded by this agreement who graduated during this period with a degree in science, mathematics, engineering, or technology fields:.....

The number of undergraduates funded by your agreement who graduated during this period and will continue to pursue a graduate or Ph.D. degree in science, mathematics, engineering, or technology fields:.....

Number of graduating undergraduates who achieved a 3.5 GPA to 4.0 (4.0 max scale):.....

Number of graduating undergraduates funded by a DoD funded Center of Excellence grant for Education, Research and Engineering:.....

The number of undergraduates funded by your agreement who graduated during this period and intend to work for the Department of Defense .....

The number of undergraduates funded by your agreement who graduated during this period and will receive scholarships or fellowships for further studies in science, mathematics, engineering or technology fields:.....

---

### Names of Personnel receiving masters degrees

NAME

**Total Number:**

---

### Names of personnel receiving PHDs

NAME

**Total Number:**

---

### Names of other research staff

NAME

PERCENT SUPPORTED

**FTE Equivalent:**

**Total Number:**

---

**Sub Contractors (DD882)**



**Inventions (DD882)**

**Scientific Progress**

**Technology Transfer**

# Fractional Differential and Integral Inequalities with Applications

---

*Donna Stutson and A. S. Vatsala*

## Table of Contents

- 1) Introduction
- 2) Preliminaries
- 3) Generalized Monotone Method for Caputo Fractional Differential Equation with Initial Condition
- 4) Generalized Monotone Method for Fractional Reaction Diffusion Equations
- 5) A Representative Formula for the Linear Caputo Fractional Wave Equation
- 6) Sub Hyperbolic Linear Partial Fractional Differential Equation in One Dimensional Space with Numerical Results
- 7) Super Hyperbolic Linear Partial Fractional Differential Equation in One Dimensional Space
- 8) Riemann Liouville and Caputo Fractional Differential and Integral Inequalities
- 9) Numerical graphs
- 10) Conclusion

## Introduction

It is well known that fractional differential equations both ordinary and fractional partial differential have played an important role in applications in theoretical modeling of scientific and engineering problems. The qualitative studies such as existence, uniqueness and computation of such dynamical models are useful in applications. Some approximate methods like the radial functions method and other methods are used in [3, 4, 7, 16] to compute exact or approximate solutions.

In solving nonlinear problems, the monotone method combined with the method of upper and lower solutions is a popular choice, because the existence of solutions by the monotone method is both theoretical and computational. The monotone method for various nonlinear problems has been developed in reference [8]. The monotone method (monotone iterative technique) combined with the method of lower and upper solutions yields monotone sequences, which converges to minimal and maximal solutions of the nonlinear differential equation. In many nonlinear problems (nonlinear dynamic systems), the nonlinear term is the sum of increasing and decreasing functions. The monotone method extended to such systems is called the generalized monotone method. Here the generalized monotone method has been extended to the Caputo fractional differential equation of order  $q$  (where  $0 < q < 1$ ) with an initial condition as well as the existence of coupled minimal and maximal solutions for such an equation and a numerical example is provided as an application of the theoretical results.

The linear sub hyperbolic fractional partial differential equation in one dimensional space, that is the  $q$ th time order derivative is such that  $1 < q < 2$ , and the linear super hyperbolic fractional partial differential equation in one dimensional space, that is the  $q$ th time order derivative is such that  $2 < q < 3$  are considered. In

the special case when  $q = 1$ , it is the parabolic equation and when  $q = 2$ , it is the hyperbolic equation. The eigenfunction expansion method is used to obtain representation forms for the solution of the linear sub and the linear super hyperbolic fractional partial differential equations in terms of the nonhomogeneous terms, initial and boundary conditions. See [9,10, 11, 12, 14] and the references therein for qualitative study, applications and numerical study of fractional differential equations. The representation forms obtained are useful in computing numerical solutions to the sub and super hyperbolic partial differential equations.

## Preliminaries

In this section, we recall some known definitions and known results, which are useful to develop our main results. Initially, we recall some definitions:

**Definition 2.1.** *Caputo derivative of order  $q$ , when  $N - 1 < q < N$  for  $t \in [0, T]$ , is defined as*

$${}^c D_{0+}^q u(t) = \frac{1}{\Gamma(N-q)} \left( \int_0^t (t-s)^{N-q-1} u^{(N)}(s) ds \right).$$

In the special case when  $N = 1$ ,

**Definition 2.2.** *Caputo fractional integral of order  $q$ , when  $0 < q < 1$ , is defined as*

$${}^c D_{0+}^q u(t) = \frac{1}{\Gamma(1-q)} \left( \int_0^t (t-s)^{-q} u'(s) ds \right).$$

**Definition 2.3.** *Riemann Liouville fractional derivative of order  $q$ , when  $0 < q < 1$  for  $t \in [0, T]$ , is defined as*

$$D^q u(t) = \frac{1}{\Gamma(1-q)} \frac{d}{dt} \int_0^t (t-s)^{-q} u(s) ds.$$

**Definition 2.4.** *Riemann Liouville right-fractional integral of order  $q$ , when  $0 < q < 1$  for  $t \in [0, T]$ , is defined as*

$${}_{t_0} D^{-q} u(t) = \frac{1}{\Gamma(q)} \frac{d}{dt} \int_0^t (t-s)^{-q} u(s) ds.$$

**Definition 2.5.** *The two parameter Mittag-Leffler function is defined as*

$$E_{\alpha, \beta}(\lambda t^\alpha) = \sum_{k=0}^{\infty} \frac{(t^\alpha)^k}{\Gamma(\alpha k + \beta)}$$

and the single parameter Mittag-Leffler function is defined as  $E_\alpha(\lambda t^\alpha) = E_{\alpha, 1}(\lambda t^\alpha)$ .

**Definition 2.6.** *let  $0 < q < 1$ ,  $p = 1 - q$  a function  $\phi \in C(J, R)$  is a  $C_p$  continuous function if  $(t - t_0)^{1-q} \phi(t) \in C(J_0, R)$ . The set of  $C_p$  functions is denoted by  $C_p(J, R)$ . Furthermore, given a function  $\phi(t) \in C(J_0, R)$ , the function  $(t - t_0)^{1-q} \phi(t)$  is called the continuous extension of  $\phi(t)$ .*

Consider the linear Caputo fractional differential equation of order  $q$  when  $N - 1 < q < N$  with initial conditions of the form

$$({}^c D^q u)(t) - \lambda u(t) = f(t), u^{(k)}(0) = b_k \quad (2.1)$$

where  $b_k \in \mathbb{R}$  for  $k = 0, 1, 2, \dots, N - 1$  and  $f \in C[J, \mathbb{R}]$  and  $J = [0, T]$ . Also,  $u^{(k)}(0)$  denotes the  $k$ th derivative of  $u$  with respect to  $t$  at  $t = 0$ . If  $f(t)$  in (2.1) is in the space of  $C^{k-1}[0, T]$ . The problem (2.1) is equivalent to the Volterra integral equation

$$u(t) = \sum_{j=0}^{N-1} \frac{b_j}{j!} t^j + \frac{\lambda}{\Gamma(q)} \int_0^t \frac{u(s) ds}{(t-s)^{1-q}} + \frac{1}{\Gamma(q)} \int_0^t \frac{f(s) ds}{(t-s)^{1-q}}.$$

The solution obtained is in the space of  $C^N[0, T]$ , see [7] for details.

Consider the equation

$$(2.5) \quad ({}^c D^q u)(t) = f(t, u(t)) + g(t, u(t)), \quad u(0) = u_0, \text{ where } 0 < q < 1, \\ f(t, u(t)), g(t, u(t)) \in C[J \times R, R], f(t, u) \text{ is increasing in } u \text{ on } J \text{ and } g(t, u) \text{ is decreasing in } u \text{ on } J.$$

**Definition.** Let  $v_0, w_0 \in C^1[J, R]$ .  $v_0, w_0$  are said to be coupled lower and upper solutions of type I of (2.5) if  $({}^c D^q v)(t) \leq f(t, v(t)) + g(t, w(t))$ ,  $v(0) \leq u_0$ , and  $({}^c D^q w)(t) \geq f(t, w(t)) + g(t, v(t))$ ,  $w(0) \geq u_0$ .

## Generalized Monotone Method for Caputo Fractional Differential Equation with an Initial Condition

The qualitative study of fractional differential equations has gained significant importance due to its applications in various branches of Science, Engineering and medicine, see [1,3,13,14,16,18]. In [14] it has been demonstrated that half order fractional differential equations give considerably better results for certain electro chemical problems than the classical approach. In addition [1,13,18] have studied linear fractional reaction diffusion equations which are applicable in random walks and nanotechnology.

Consider the equation

$$(3.1) \quad ({}^c D^q u)(t) = f(t, u(t)) + g(t, u(t)), \quad u(0) = u_0, \\ \text{where } 0 < q < 1, f(t, u(t)), g(t, u(t)) \in C[J \times R, R], f(t, u) \text{ is increasing in } u \text{ on } J \text{ and } g(t, u) \text{ is decreasing in } u \text{ on } J.$$

**Theorem 3.1** Assume that

- I.  $v_0, w_0 \in C^1[J, R]$ .  $v_0, w_0$  are coupled lower and upper solutions of type I, with  $v_0 \leq w_0$  on  $J$ .
- II.  $f(t, u(t)), g(t, u(t)) \in C[J \times R, R]$ , where  $f(t, u)$  is increasing in  $u$  on  $J$  and  $g(t, u)$  is decreasing in  $u$  on  $J$ .

Then there exist monotone sequences,  $v_n(t)$  and  $w_n(t)$ , such that  $v_n(t) \rightarrow \rho(t)$  and  $w_n(t) \rightarrow r(t)$  uniformly and monotonically, where  $\rho(t)$  and  $r(t)$  are coupled minimal and maximal solutions of equation (3.1) on  $J$ . That is, for any solution,  $u(t)$ , of (3.1) with  $v_0(t) \leq u_0(t) \leq w_0(t)$  on  $J$ , we get natural sequences,  $\{v_n\}$  and  $\{w_n\}$ , satisfying the following,

$$v_0(t) \leq v_1(t) \leq v_2(t) \leq \dots \leq v_n(t) \leq u(t) \leq w_n(t) \leq \dots \leq w_2(t) \leq w_1(t) \leq w_0(t),$$

for each  $n \leq 1$  on  $J$ . Also  $\rho(t) \leq u(t) \leq r(t)$  on  $J$ .

The next result uses intertwined sequences.

**Theorem 3.2.** Assume hypotheses, I. and II, of the above theorem hold. Then for any solution,  $u(t)$ , of equation (3.1) with  $v_0(t) \leq u_0(t) \leq w_0(t)$  on  $J$ , the alternating sequences,  $\{v_{2n}, w_{2n+1}\}$  and  $\{v_{2n+1}, w_{2n}\}$ , satisfying intertwined sequences,

(3.7)  $v_0(t) \leq w_1(t) \leq \dots \leq v_{2n}(t) \leq w_{2n+1}(t) \leq u(t) \leq v_{2n+1}(t) \leq w_{2n}(t) \leq \dots \leq v_1(t) \leq w_0(t)$ , for each  $n \geq 1$  on  $J$ . This requires using type (iii) iterative schemes,

$$\begin{aligned} ({}^c D^q v_{n+1})(t) &= f(t, w_n(t)) + g(t, v_n(t)), v_{n+1}(0) = u_0 \text{ and} \\ ({}^c D^q w_{n+1})(t) &= f(t, v_n(t)) + g(t, w_n(t)), w_{n+1}(0) = u_0. \end{aligned}$$

Further, monotone sequences,  $\{v_{2n}, w_{2n+1}\}$  and  $\{w_{2n}, v_{2n+1}\}$ , converge to  $\rho(t)$  and  $r(t)$ , respectively, on  $J$ .  $\rho(t)$  and  $r(t)$  are coupled minimal and maximal solutions of (3.1), respectively. Also,  $\rho(t) \leq u(t) \leq r(t)$  on  $J$ .

**Theorem 3.4.** If in addition to hypotheses I and II, suppose  $f(t, u)$  satisfies left-hand-side Lipschitz condition,  $f(t, u_2) - f(t, u_1) \leq L_1(u_2 - u_1)$ ,  $u_2 \geq u_1$ , and  $g(t, u)$  satisfies right-hand-side Lipschitz condition,  $g(t, u_2) - g(t, u_1) \geq -L_2(u_2 - u_1)$ ,  $u_2 \geq u_1$ , where  $v_0 \leq u_1 \leq u_2 \leq w_0$  implies that  $\rho = r = u$ , the unique solution of (3.1).

## GENERALIZED MONOTONE METHOD FOR FRACTIONAL REACTION DIFFUSION EQUATIONS

Consider the fractional reaction diffusion equation of the form

$$\begin{aligned} {}^c \partial_t^q u - k \frac{\partial^2 u}{\partial x^2} &= f(t, x, u) + g(t, x, u), \quad (t, x) \in Q_T \\ u(t, 0) &= A(t), \quad u(t, L) = B(t) \quad (t, x) \in \Gamma_T \\ u(0, x) &= h(x) \quad x \in \bar{\Omega} = \Omega, \end{aligned} \tag{4.1}$$

where  $\Omega = [0, L]$ ,  $J = (0, T]$ ,  $Q_T = J \times \Omega$ ,  $k > 0$  and  $\Gamma_T = (0, T) \times \partial\Omega$ .

Here  ${}^c \partial_t^q u$  is the Caputo Partial Derivative with respect to  $t$  of order  $q$ ,  $0 < q < 1$ , which is defined as

$${}^c \partial_t^q u = \frac{1}{\Gamma(1-q)} \left( \int_0^t (t-s)^{-q} \frac{\partial u(s, x)}{\partial s} ds \right).$$

Assume that  $f, g \in C^{1,2}[Q_T \times R, R]$ , such that  $f(t, x, u)$  is non-decreasing in  $u$  and  $g(t, x, u)$  is non-increasing in  $u$ ,  $A(t), B(t) \in C^1[J, R]$  and  $h(x) \in C^{2+\alpha}[\Omega, R]$ , where  $0 < \alpha < 1$ .

In this section we develop the generalized monotone method for (4.1) combined with coupled lower and upper solutions of type I. The generalized monotone method yields monotone sequences which converge uniformly and monotonically to coupled minimal and maximal solutions of (4.1). Furthermore, assuming a uniqueness condition, we prove the existence of the unique solution of (4.1).

**Theorem 4.1:** Let  $(v_0, w_0)$  be coupled lower and upper solutions of (2.1) such that  $v_0 \leq w_0$  on  $Q_T$ . Furthermore, suppose  $f(t, x, u)$  is nondecreasing in  $u$  on  $Q_T$  and  $g(t, x, u)$  is nonincreasing in  $Q_T$  and

$$\begin{aligned} f(t, x, u) - f(t, x, \bar{u}) &\leq L_1(u - \bar{u}) \\ g(t, x, u) - g(t, x, \bar{u}) &\geq -L_2(u - \bar{u}), \end{aligned}$$

whenever  $u \geq \bar{u}$  on  $Q_T$ . Then there exists monotone sequences  $\{v_n(t, x)\}$  and  $\{w_n(t, x)\}$  defined such that  $v_n(t, x)$  and  $w_n(t, x)$  are solutions to

$$\begin{aligned} {}^c\partial_t^q v_n(t, x) - k \frac{\partial^2 v_n(t, x)}{\partial x^2} &= f(t, x, v_{n-1}) + g(t, x, w_{n-1}), \quad (t, x) \in Q_T \\ v_n(t, 0) &= A(t), \quad v_n(t, L) = B(t) \quad (t, x) \in \Gamma_T \\ v_n(0, x) &= h(x) \quad x \in \bar{\Omega} = \Omega. \end{aligned} \quad (4.2)$$

and

$$\begin{aligned} {}^c\partial_t^q w_n(t, x) - k \frac{\partial^2 w_n(t, x)}{\partial x^2} &= f(t, x, w_{n-1}) + g(t, x, v_{n-1}), \quad (t, x) \in Q_T \\ w_n(t, 0) &= A(t), \quad w_n(t, L) = B(t) \quad (t, x) \in \Gamma_T \\ w_n(0, x) &= h(x) \quad x \in \bar{\Omega} = \Omega. \end{aligned} \quad (4.3)$$

respectively. Also, these sequences converge uniformly and monotonically to  $\rho(t, x)$  and  $r(t, x)$  such that  $(\rho(t, x), r(t, x))$  are coupled minimal and maximal solutions of the nonlinear initial value problem (4.1), provided  $v_0(t, x) \leq u(t, x) \leq w_0(t, x)$ , where  $u(t, x)$  is any solution of (2.1). Furthermore,  $\rho(t, x) = r(t, x) = u(t, x)$ , the unique solution of (2.1), on  $\bar{Q}_T$ .

## A Representative Formula for the Linear Caputo Fractional Wave Equation

It is well known that fractional Brownian motion has been modeled as parabolic stochastic differential equation. Here a representation form for the solution of the deterministic one dimensional fractional wave equation with Caputo fractional derivative of order  $q$ , for  $1 < q < 2$  has been developed. For  $q = 1$  and  $q = 2$ , it reduces to the one dimensional parabolic equation and one dimensional wave equation, respectively.

Consider the linear Caputo fractional wave differential equation

$$\begin{aligned} {}^c\partial_t^q u - c^2 \frac{\partial^2 u}{\partial x^2} &= R(t, x), \quad (t, x) \in Q_T \\ u(t, 0) &= A(t), \quad u(t, L) = B(t) \quad (t, x) \in \Gamma_T \\ u(0, x) &= h(x), u_t(0, x) = g(x) \quad x \in \bar{\Omega}, \end{aligned} \quad (5.1)$$

Where  $\Omega = [0, L], J = (0, T], Q_T = J \times \Omega, c^2 > 0$  and  $\Gamma_T = (0, T) \times \partial\Omega$ .

In order to obtain a representation formula we need to consider the corresponding homogeneous equation with homogeneous boundary condition. Using separation of variables method we can find the eigenfunctions corresponding to the eigenvalues  $\lambda_n$ . The eigenfunctions and the eigenvalues vary depending on the type of

boundary conditions. In order to get a suitable final form of the solution we normalize the eigenfunctions. For the above type of boundary conditions, we get  $\lambda_n = \left(\frac{n\pi}{L}\right)^2$  and the eigenfunctions are  $\phi_n(x) = \sin \frac{n\pi x}{L}$ .

$$u(t, x) = \int_0^L \left[ \sum_{n=1}^{\infty} h(x_0) \phi_n(x_0) dx_0 E_{q,1}(-c^2 \lambda_n t^q) + \int_0^L \left[ \sum_{n=1}^{\infty} g(x_0) \phi_n(x_0) dx_0 t^q E_{q,2}(-c^2 \lambda_n t^q) \right. \right. \\ \left. \left. + \int_0^t (t-s)^{q-1} E_{qq}(-c^2 \lambda (t-s)^q) \int_0^L Q(s, x) ds + \frac{c^2 n\pi}{L} \int_0^t (t-s)^{q-1} E_{qq}(-c^2 \lambda (t-s)^q) A(s) ds \right. \right. \\ \left. \left. + (-1)^{n+1} \frac{c^2 n\pi}{L} \int_0^t (t-s)^{q-1} E_{qq}(-c^2 \lambda (t-s)^q) B(s) ds \right] \right].$$

## Sub Hyperbolic Linear Partial Fractional Differential Equation in One Dimensional Space with Numerical Results

In this section a representation form for the sub hyperbolic linear fractional partial differential equation in one dimensional space is obtained. It is called sub hyperbolic if the time derivative is of order  $q$  when  $1 < q < 2$ . It is easy to observe that, if  $q = 2$ , it is a second order linear hyperbolic equation in one dimensional space. Consider the following linear Caputo fractional wave differential equation:

$$\begin{aligned} {}^c \partial_t^q u - c^2 \frac{\partial^2 u}{\partial x^2} &= F(x, t) \text{ on } Q_T, \\ u(0, t) &= A(t), \quad u(L, t) = B(t) \quad \text{on } \Gamma_T, \\ u(0, x) &= h_0(x), \quad u_t(0, x) = h_1(x) \quad \text{on } \bar{\Omega}, \end{aligned} \quad (6.1)$$

where  $\Omega = [0, L]$ ,  $J = (0, T]$ ,  $Q_T = J \times \Omega$ ,  $c^2 > 0$  and  $\Gamma_T = (0, T) \times \partial\Omega$ .

We develop a representation form for (6.1) which is useful in computing the solution of linear Caputo fractional partial differential equations of order  $q$  when  $1 < q < 2$ , using the eigenfunction expansion method with eigenvalues  $\lambda_n = \left(\frac{n\pi}{L}\right)^2$  and eigenfunctions  $\phi_n(x) = \sin\left(\frac{n\pi x}{L}\right)$ . This will be useful in the development of the generalized monotone method to be developed in future work. We also assume that the initial boundary condition satisfy the compatability condition. Hence, we set

$$u(t, x) = \sum_{n=1}^{\infty} b_n(t) \sin \frac{n\pi x}{L}. \quad (6.2)$$

We must then find  $b_n(t)$  for each  $n \geq 1$ . Taking the Caputo derivative of each side, Green's formula and the boundary conditions yields the following fractional equation to be solved:

$${}^c D^q b_n(t) + c^2 \left(\frac{n\pi}{L}\right)^2 b_n(t) = \frac{2}{L} \left[ \frac{n\pi c^2}{L} \{A(t) + (-1)^n B(t)\} + \int_0^L F(t, x) \sin\left(\frac{n\pi x}{L}\right) dx \right] \quad (6.3)$$

With initial conditions

$$b_n(0) = \frac{2}{L} \int_0^L h_0(x) \sin\left(\frac{n\pi x}{L}\right) dx, \quad b_n'(0) = \frac{2}{L} \int_0^L h_1(x) \sin\left(\frac{n\pi x}{L}\right) dx.$$

Using Equation (2.1) to solve (6.3) substituting in for  $b_n(t)$  in equation (6.2) and interchanging the order of integration and summation, we obtain the representation form:

$$u(t, x) =$$

$$\begin{aligned}
& \int_0^L \left[ \sum_{n=1}^{\infty} h(x_0) \varphi_n(x_0) dx_0 E_{q,1}(-c^2 \lambda_n t^q) \right] \\
& + \int_0^L \left[ \sum_{n=1}^{\infty} h_1(x_0) \varphi_n(x_0) \varphi_n(x) dx_0 t E_{q,2}(-c^2 \lambda_n t^q) \right] \\
& + \frac{2}{L} \int_0^t \int_0^L \sum_{n=1}^{\infty} (t-s)^{(q-1)} E_{q,q}(-c^2 \lambda_n (t-s)^q) [F(s, x_0) \varphi_n(x_0) \varphi_n(x) dx_0] ds \\
& + \int_0^t \frac{2c^2 n \pi}{L^2} (t-s)^{(q-1)} E_{q,q}(-c^2 \lambda_n (t-s)^q) A(s) \varphi_n(x) ds \\
& + (-1)^{n+1} \int_0^t \frac{2c^2 n \pi}{L^2} (t-s)^{(q-1)} E_{q,q}(-c^2 \lambda_n (t-s)^q) B(s) \varphi_n(x) ds.
\end{aligned}$$

## Super Hyperbolic Linear Partial Fractional Differential Equation in One Dimensional Space with Numerical Results

Using the method of successive approximation by setting  $u_0(t) = \sum_{j=0}^{N-1} \frac{b_j}{j!} t^j$  and taking the limit of  $u_m(t)$  as  $m \rightarrow \infty$ , the solution of (2.1) is given by

$$u(t) = \sum_{j=0}^{N-1} b_j t^j E_{q,j+1}(\lambda t^q) + \int_0^t (t-s)^{q-1} E_{q,q}(\lambda(t-s)^q) f(s) ds.$$

Next we consider an eigenvalue problem on one dimensional space. For that purpose consider the eigenvalue problem

$$u'' + \lambda u = 0$$

with mixed homogeneous boundary conditions of the form:

$$B_u(a) = \alpha_i u(i) + (-1)^{(i+1)} \beta_i u'(i) = 0, \text{ for } i = 0, 1.$$

The value of  $\lambda$  for which the nontrivial solution for the boundary value problem exists is called the eigenvalue denoted as  $\lambda_n$  and the corresponding nontrivial solution is called the eigenfunction denoted as  $\varphi_n(x)$ . The main result is developed using the eigenvalue problem with Dirichlet boundary conditions,

$$u(0) = u(L) = 0.$$

In this section a representation form for the super hyperbolic linear fractional partial differential equation in one dimensional space is obtained. It is called super hyperbolic if the time derivative is of order  $q$  when  $2 < q < 3$ . It is easy to observe that, if  $q = 2$ , it is a second order linear hyperbolic equation in one dimensional space. Consider the following linear Caputo fractional wave differential equation:

$$\begin{aligned}
& {}^c \partial_t^q u - c^2 \frac{\partial^2 u}{\partial x^2} = F(x, t) \text{ on } Q_T, \\
& u(0, t) = A(t), \quad u(L, t) = B(t) \quad \text{on } \Gamma_T, \\
& u(0, x) = h_0(x), \quad u_t(0, x) = h_1(x), \quad u_{tt}(0, x) = h_2(x) \quad \text{on } \bar{\Omega},
\end{aligned} \tag{7.1}$$

where  $\Omega = [0, L]$ ,  $J = (0, T]$ ,  $Q_T = J \times \Omega$ ,  $c_2 > 0$  and  $\Gamma_T = (0, T) \times \partial\Omega$ . We also assume that  $A(t), B(t) \in C^3[J, R]$ , and  $F(t, x) \in C^{3,2}[Q_T, R]$ .

Assume the following compatibility conditions relative to the initial and boundary conditions:

$$h_0(0) = A(0) = 0, \quad h_0(L) = B(0) = 0$$



$$h_1(0) = A'(0) = 0, \quad h_1(L) = B''(0) = 0$$

$$h_2(0) = A''(0) = 0, \quad h_2(L) = B''(0) = 0.$$

This is needed when differentiating the integral representation of the terms related to the boundary conditions and when the problem is reduced to homogeneous boundary conditions. As in section 3, we set  $u(t, x) = \sum_{n=1}^{\infty} b_n(t) \sin \frac{n\pi x}{L}$ . As before, the expression for  $b_n(t)$  is found for each  $n \geq 1$  by using Lagrange's identity along with Green's formula and the boundary conditions and substitute back into the expression for  $u(t, x)$ . Changing the order of integration and summation, we obtain the solution :

$$u(t, x) =$$

$$\begin{aligned} & \int_0^L \left[ \sum_{n=1}^{\infty} \sum_{j=0}^2 h_j(x_0) \varphi_n(x_0) dx_0 t^j E_{q,j+1}(-c^2 \lambda_n t^q) \right] \\ & + \frac{2}{L} \int_0^t \int_0^L \sum_{n=1}^{\infty} (t-s)^{(q-1)} E_{q,q}(-c^2 \lambda_n (t-s)^q) [F(s, x_0) \varphi_n(x_0) \varphi_n(x) dx_0] ds \\ & + \int_0^t \sum_{n=1}^{\infty} \frac{2c^2 n\pi}{L^2} (t-s)^{(q-1)} E_{q,q}(-c^2 \lambda_n (t-s)^q) A(s) \varphi_n(x) ds \\ & + (-1)^{n+1} \int_0^t \frac{2c^2 n\pi}{L^2} (t-s)^{(q-1)} E_{q,q}(-c^2 \lambda_n (t-s)^q) B(s) \varphi_n(x) ds. \end{aligned}$$

This representation formula for the solution of (7.1) provides a methodology for computing the solution for different nonhomogeneous terms and different initial and boundary conditions. The convergence of the infinite series involves the Mittag-Leffler functions is yet to be established even when  $0 < q < 1$ . However, if the boundary conditions are homogeneous and the initial and nonhomogeneous terms are chosen such that there are a finite number of Mittag-Leffler functions, then the solution can be numerically computed using MAPLE.

## Riemann Liouville and Caputo Fractional Differential and Integral Inequalities

In this section, results for coupled fractional and ordinary integral inequalities where the nonlinear function is the sum of an increasing and a decreasing function have been developed. Also, the corresponding coupled fractional and ordinary differential inequality results have been developed without requiring the increasing or decreasing nature of the nonlinear function. In fact, results have been developed for coupled differential and integral inequalities for both ordinary and fractional equations. These results are useful in the qualitative study of ordinary and fractional dynamic systems of both Riemann Liouville and Caputo forms.

For the sake of simplicity, the obtained results will be on the interval  $J = (t_0, T]$ . Furthermore,  $J_0 = (t_0, T] = \bar{J}$ .

**Theorem 8.1.** Let  $v, w \in C_p[J, R]$ ,  $f, g \in C[J_0 \times R, R]$  and satisfy the following coupled integral inequalities

$$(i) \ v(t) \leq \frac{v_0(t-t_0)^{q-1}}{\Gamma(q)} + \frac{1}{\Gamma(q)} \int_{t_0}^t (t-s)^{q-1} \left( f(s, v(s)) + g(s, w(s)) \right) ds,$$

$$(ii) w(t) \geq \frac{w_0(t-t_0)^{q-1}}{\Gamma(q)} + \frac{1}{\Gamma(q)} \int_{t_0}^t (t-s)^{q-1} (f(s, w(s)) + g(s, v(s))) ds,$$

(iii)  $f(t, u)$  is nondecreasing in  $u$  for each  $t \in J_0$ , and  $f(t, u)$  satisfies the one sided Lipschitz condition of the form

$$f(t, u_1) - f(t, u_2) \leq L(u_1 - u_2),$$

for some  $L > 0$ , whenever  $u_1 \geq u_2$ ;

(iv)  $g(t, u)$  is non increasing in  $u$  for each  $t \in J_0$  and  $g(t, u)$  satisfies the one sided Lipschitz condition of the form

$$g(t, u_1) - g(t, u_2) \geq -M(u_1 - u_2),$$

for some  $M > 0$ , whenever  $u_1 \geq u_2$ . Then  $u_0 \leq w_0$  implies  $v(t) \leq w(t)$  on  $J$ .

Consider the Riemann-Liouville fractional differential equation of the form:

$$D^q u(t) = f(t, u(t)) + g(t, u(t)), \quad \Gamma(q)(t - t_0)^{1-q} u(t)|_{t=t_0} = u^0, \quad (8.1)$$

where  $f, g \in C[J_0 \times R, R]$ .

The next result is the differential inequality result relative to the Riemann-Liouville fractional differential equation (8.1).

**Theorem 8.2.** Let  $v, w \in C_p[J, R]$ ,  $f, g \in C[J_0 \times R, R]$  and satisfy the following coupled differential inequalities

$$(i) D^q(v(t)) \leq f(t, v(t)) + g(t, w(t)), \quad \Gamma(q)(t - t_0)^{1-q} v(t)|_{t=t_0} \leq u^0;$$

$$(ii) D^q(w(t)) \geq f(t, w(t)) + g(t, v(t)), \quad \Gamma(q)(t - t_0)^{1-q} w(t)|_{t=t_0} \geq u^0;$$

(iii)  $f(t, u)$  satisfies the one sided Lipschitz condition of the form

$$f(t, u_1) - f(t, u_2) \leq L(u_1 - u_2),$$

for some  $L > 0$ , whenever  $u_1 \geq u_2$ ;

(iv)  $g(t, u)$  satisfies the one sided Lipschitz condition of the form

$$g(t, u_1) - g(t, u_2) \geq -M(u_1 - u_2),$$

for some  $M > 0$ , whenever  $u_1 \geq u_2$ . Then  $v_0 \leq w_0$  implies  $v(t) \leq w(t)$  on  $J$ .

Here the Caputo fractional differential and integral inequalities are developed.

Consider the Caputo fractional differential equation of order  $q$  where  $0 < q < 1$ , of the form :

$${}^C D^q u(t) = \lambda u(t) + f(t), \quad u(t_0) = u_0, \quad (8.2)$$

where  $f, g \in C[J_0 \times R, R]$ .

The integral representation of (8.2) is given by equation

$$u(t) = u_0 + E_{q,1}(\lambda(t - t_0)^q) + \int_{t_0}^t (t - s)^{q-1} E_{q,q}(\lambda(s - t_0)^q) f(s) ds, \quad (8.3)$$

where  $\Gamma(q)$  is the Gamma function. Note, the above holds true if in place of equality we have less than or equal.

Now consider the Caputo fractional differential equation

$${}^c D^q u(t) = f(t, u) + g(t, u), \quad u(t_0) = u_0, \quad (8.4)$$

Where  $f, g \in C[J_0 \times R, R]$ .

**Definition 8.1.** The functions  $v, w \in C^1([t_0, T], R)$  are called natural lower and upper solutions of (8.4) if :

$${}^c D^q v(t) \leq f(t, v) + g(t, v), \quad v(t_0) \leq u_0,$$

and

$${}^c D^q w(t) \geq f(t, w) + g(t, w), \quad w(t_0) \geq u_0.$$

**Definition 8.2.** The functions  $v, w \in C^1([t_0, T], R)$  are called coupled lower and upper solutions of type I of (8.4) if :

$${}^c D^q v(t) \leq f(t, v) + g(t, w), \quad v(t_0) \leq u_0,$$

and

$${}^c D^q w(t) \geq f(t, w) + g(t, v), \quad w(t_0) \geq u_0.$$

One can easily prove the existence of the solution of (8.4) on the interval  $[t_0, T]$  when natural lower and upper solution for (8.4) are known for which  $v(t) \leq w(t)$ . Coupled minimal and maximal solutions of (8.4) on the interval  $[t_0, T]$  can be computed without any extra assumptions using the generalized monotone method when coupled lower and upper solutions of type I of (8.4) are known with  $v(t) \leq w(t)$ .

Next we have the differential inequality result for coupled lower and upper solutions of type I:

**Theorem 8.3.** Let  $v, w \in C^1(J, R)$ , Where  $f, g \in C[J \times R, R]$  are coupled lower and upper solutions of type I, such that

(i)  $f(t, u)$  satisfies the one sided Lipschitz condition of the form

$$f(t, u_1) - f(t, u_2) \leq L(u_1 - u_2),$$

for some  $L > 0$ , whenever  $u_1 \geq u_2$ ;

(ii)  $g(t, u)$  satisfies the one sided Lipschitz condition of the form

$$g(t, u_1) - g(t, u_2) \geq -M(u_1 - u_2),$$

for some  $M > 0$ , whenever  $u_1 \geq u_2$ . Then  $v(t_0) \leq w(t_0)$  implies that  $v(t) \leq w(t)$ , on  $J$ .

**Remark 5.1.** Caputo integral inequalities results similar to Theorem 8.1 can also be developed. The integral inequality results need an extra assumption which is not required for the corresponding scalar differential inequality results. Thus, the differential inequality results are useful in iterative methods like the monotone quasilinearization methods. They are useful for proving uniqueness, continuous dependency on the initial condition and also in determining the order of convergence.

## Numerical Graphs

Consider a special case of the Logistic model of the form

$$(4.1) \quad {}^c D^{\frac{1}{2}} u(t) = 0.99u(t) - u^2(t), \quad u(0) = \frac{1}{2}.$$




FIGURE 2. Dashed:  $\{v_n\}$ . Solid:  $\{w_n\}$ .

## Conclusion

We have developed generalized monotone method for nonlinear fractional differential systems and fractional reaction diffusion equations via coupled lower and upper solutions of type 1. The advantage of the generalized monotone method is that each component of the iterates are scalar fractional differential equation whose solutions are easy to compute compared to computing the solution corresponding linear system. Also, This representation formulas provide a methodology for computing solutions for different nonhomogeneous terms and different initial and boundary conditions.

## References

- [1] D. Baleanu, Z. B. Guvencs, J.A. T. Machado, New Trends in Nanotechnology and Fractional Calculus Applications, Springer, 2010.
- [2] M. Caputo, Linear models of dissipation whose Q is almost independent, II, *Geophysical Journal of the Royal Astronomical Society*, **13** (1967), 529-539.
- [3] Z. Denton and A.S.Vatsala, "Monotone Iterative Technique for Finite Systems of Nonlinear Fractional Differential Equations," *Opuscula Mathematica*, Vol. 31, No.3, 2011.
- [4] Z. Denton and A.S.Vatsala, "Fractional Integral Inequalities and Applications," *Computers and Mathematics with Applications*, 59 (2010), 1087-1094.
- [5] Z. Denton, P.W. Ng, and A.S. Vatsala, Quasilinearization Method via Lower and Upper Solutions for Riemann-Liouville Fractional Differential Equations, *Nonlinear Dynamics and Systems Theory*, 11 (3) (2011), 239-251.
- [6] Kai Diethelm, *The Analysis of Fractional Differential Equations*, Springer, 2004.
- [7] A. Kilbas, H. M. Srivastava, J. T. Trujillo, Theory and Applications of Fractional Differential Equations, North-Holland Mathematical studies, 204, Elsevier, 2006.

- [8] G.S. Ladde, V. Lakshmikantham and A.S. Vatsala, Monotone Iterative Techniques for Nonlinear Differential Equations, Pitman Publishing, Inc., 1985.
- [9] V. Lakshmikantham and A. S. Vatsala, *Generalized Quasilinearization for Nonlinear Problems*, Kluwer Academic Publishers, Dordrecht, Boston, London.
- [10] V. Lakshmikantham, S. Leela and J. Vasundara Devi, Theory of Fractional Dynamic Systems, Cambridge Scientific Publishers, Ltd, 2009.
- [11] G. S. Ladde, V. Lakshmikantham and A.S. Vatsala, *Monotone Iterative Techniques for Nonlinear Differential Equations*, Pitman (1985).
- [12] V. Lakshmikantham, S. Leela, *Differential and Integral Inequalities*, Volume I, II, Academic Press, New York, 1969.
- [13] R. Metzler and J. Klafter, "The random walks's guide to anomalous diffusion: a fractional dynamics approach," *Phys Rep* 339(1):1-77.
- [14] B. Oldham and J. Spanier, The Fractional Calculus, Academic Press, New York, London, 1974.
- [15] TH.T. Pham. J. D. Ramirez, A. S. Vatsala, Generalized Monotone Method for Caputo Fractional Differential Equations with Applications to Population Models, *Neural, Parallel and Scientific Computations*, 20(2012), 119-132.
- [16] I. Podlubny, Fractional Differentials Equations, Mathematics in Science and Engineering, Volume 198, Academic Press, 1999.
- [17] J. D. Ramirez, A.S. Vatsala, "Generalized Monotone Iterative Technique for Caputo Fractional Differential Equation with Periodic Boundary Condition via Initial Value Problem," submitted for publication to *Integral Inequalities and Applications*.
- [18] M. Rivero, J. Trujillo and M. Pilar Velasco, "On Deterministic Fractional Models," New Trends in Nanotechnology and Fractional Calculus Applications, edited by D. Baleanu, et al, 123-150, Springer, 2010.
- [19] M. Sokol and A.S. Vatsala, "A unified exhaustive study of monotone iterative method for initial value problems," *Nonlinear Studies*, 8 (4), 429-438,(2001).
- [20] M. Sowmya and A. S. Vatsala, Numerical Approach via Generalized Monotone Method for Scalar Caputo Fractional Differential Equations, *Neural, Parallel and Scientific Computations* 21(2013), 19-30.
- [21] M. Sowmya and A. S. Vatsala, Superlinear Convergence for Caputo Fractional Differential Equations with Applications, *Dynamic Systems and Applications* 22(2013), 479-492.
- [22] D. Stutson and A. S. Vatsala, Generalized Monotone Method for Caputo Fractional Differential Systems via Coupled Lower and Upper Solutions, *Dynamic Systems and Applications* 20(2011), 495-504.
- [23] A.S. Vatsala and I. H. West, "Generalized Monotone Iterative Method for Initial Value Problems," *Applied Mathematics Letters*, 17 (2004) 1231-1237.